

# ANTIOXIDANT STATUS IN ASYMPTOMATIC HYPERCHOLESTEROLEMIC SUBJECTS

Simmi Kharb

Department of Biochemistry, Pandit Bhagwat Dayal Sharma Postgraduate Institute of Medical Sciences, Rohtak, India

## ABSTRACT

**Background:** An imbalance between antioxidant and oxidant-generating systems leading to an oxidative stress has already been proposed in the pathogenesis of atherosclerosis. In the present study, the antioxidant status in asymptomatic hypercholesterolemic persons was compared with normocholesterolemics.

**Methodology:** In this study, the antioxidant status in 30 asymptomatic hypercholesterolemic persons was determined and compared with 15 normocholesterolemic ones.

**Results:** Hypercholesterolemics had significantly lower vitamin E, superoxide dismutase and glutathione levels as compared to normocholesterolemics. Serum lipid peroxidation product, malonaldehyde levels were significantly higher in Hypercholesterolemics.

**Conclusion:** These observations demonstrate altered antioxidant status in asymptomatic hypercholesterolemic persons. Low vitamin E detected in hypercholesterolemics supports the hypothesis that vitamin E deficiency manifests in the early stage of atherosclerosis and might be the most sensitive parameter for this.

**KEYWORDS:** Antioxidant, Atherosclerosis, Cholesterol, vitamin E.

## INTRODUCTION

Accumulating evidence suggests that alterations or disturbances in defensive antioxidant system may contribute to or exacerbate the development of atherosclerosis.<sup>1,2</sup> The so-called "oxidation hypothesis" states that the oxidative modification of low density lipoprotein (LDL) or other lipoproteins is important and possibly obligatory in the pathogenesis of the atherosclerotic lesions.<sup>3,4</sup>

The antioxidant defenses that prevent oxidation of LDL need to be defined. The antioxidant content of the LDL particle is critical for its protection<sup>5</sup> and the most abundant natural antioxidant in LDL is alpha-tocopherol.<sup>6</sup> The potential antiatherogenic role of vitamin E has been evaluated from epidemiological studies.<sup>7</sup> It has been reported that in hypercholesterolemia, erythrocyte vitamin E content was decreased although their plasma vitamin E concentration was unchanged compared with normocholesterolemic men.<sup>8</sup> Also, erythrocyte enzymes superoxide dismutase (SOD) and glutathione peroxidase (GPx) contribute to the cell antioxidant defense mechanisms. Decrease in SOD and/or GPx activities has been described in stable angina and coronary heart disease.<sup>8</sup> While increased SOD was found in unstable angina reflecting an adaptive induction of the enzyme in order to limit the excess free radical production.<sup>9</sup>

All these results are available for subjects with symptomatic cardiovascular diseases, but few studies investigated the antioxidant status in a stage of preclinical disease.

The present study was aimed to investigate the antioxidant status in subjects free from any symptomatic cardiovascular diseases, but with a well-established cardiovascular risk factor, hypercholesterolemia (HC), compared with normocholesterolemic (NC) subjects.

## MATERIAL AND METHODS

This study was carried out in 30 HC subjects, (20 M, 10 F), with a total cholesterol level >240 mg/dl and they were compared with 15 NC (11 M, 6 F) subjects.

All the subjects were free of secondary hypercholesterolemia, hypertriglyceridemia (TG >175 mg/dl), renal failure, diabetes mellitus or history of myocardial infarction, stroke or intermittent claudication and were not taking any vitamin supplements or lipid-lowering drugs.

Overnight fasting venous samples were drawn. Serum total cholesterol and triglycerides were determined enzymatically. Also, serum malonaldehyde (MDA),<sup>10</sup> and vitamin E<sup>11</sup> were analyzed. SOD activity was measured by Misra and

Fridovich method.<sup>12</sup> Glutathione levels were analyzed by Beutler's method.<sup>13</sup> The vitamin E total lipid ratio was calculated.

The data obtained was analyzed statistically using student's t-test.

## RESULTS

Serum lipid peroxidation product MDA levels were significantly higher in HC as compared to NC subjects, with  $p < 0.05$ .

On the other hand, antioxidant defenses: vitamin E, SOD and GSH were significantly lower in HC as compared to NC subjects.

Vitamin E: total lipids ratio was lower in HC as compared to NC. (Table 1)

**Table 1: Parameters of oxidative stress (mean+SD).**

	<b>Hypercholesterolemic persons (n=30)</b>	<b>Normocholesterolemic persons (n=15)</b>
Vitamin E ( $\mu\text{mol/L}$ )	19.93 + 2.51	25.75 + 2.89
SOD(EU/L)	173 + 2.51	256.12 + 6.09
MDA (nmol/ml)	0.86 + 0.27	0.69 + 0.02
GSH(mg/dl)	35.5 + 6.8	42.3 + 11.6*
Vitamin E/ Total lipids ( $\mu\text{mol}/\text{mmol}$ )	6.94 + 0.17	8.67 + 6.26

Values are significant at all levels ( $p < 0.05$ ) except\*

## DISCUSSION

The present study was designed to assess the antioxidant status in subjects free from asymptomatic cardiovascular disease, but with a well-established cardiovascular risk factor, hypercholesterolemia, compared with normocholesterolemic subjects. The vitamin E concentrations were significantly lower in HC compared with NC. Our findings are consistent with those reported in the literature.<sup>8</sup> Vitamin E localizes mainly to lipoproteins and membranes, where it serves to scavenge lipid peroxyl radicals.<sup>6</sup> Despite this activity, atherosclerosis is characterized by lipid peroxidation within the vascular wall even in the presence of alpha-tocopherol independent of its effect to inhibit lipid peroxidation. This is shown in a report, where depleting the diet of total toco-

pherols in rabbits resulted in a dose-dependent impairment of NO-mediated relaxation upon exposure to OX-LDL.<sup>15</sup> In contrast, vessels containing abundant alpha-tocopherol content were markedly resistant this effect of OX-LDL.<sup>15</sup> One hallmark of the atherosclerotic plaque is proliferation of vascular smooth muscles. Patients who demonstrate rapid lesion progression also exhibit excess cardiovascular events.<sup>15</sup> Considerable in vitro data indicate that vitamin E inhibits the proliferation of smooth muscle cells.<sup>16</sup> And reports suggest that alpha-tocopherol inhibits smooth muscle cell proliferation principally as a consequence of protein kinase C inhibition.<sup>16</sup> In the setting of atherosclerosis, inhibition of protein kinase C by vitamin E would be expected to maintain normal vascular homeostasis and thus reduce the clinical incidence of cardiovascular disease.

Also, SOD activities were significantly lower in HC compared with NC and GSH content was lower in HC, but, it was statistically insignificant. Several studies have indicated that superoxide production is elevated in atherosclerotic vessels.<sup>17</sup> This increased production of superoxide has been shown to interfere with EDRF-dependent relaxations of vessels.<sup>17</sup> Intriguingly, administration of Cu Zn-SOD encapsulated in liposomes results in an increased level of SOD activity in the extracellular matrix.<sup>17</sup> This elevation in SOD activity augments EDRF-dependent relaxation in atheromatous vessels. Finding of low GSH levels in HC in addition to elevated lipid peroxidation products suggest existence of oxidative stress in HC subjects.

## CONCLUSION

These results demonstrate altered antioxidant status in asymptomatic hypercholesterolemic persons. Low vitamin E detected in hypercholesterolemics supports the hypothesis that vitamin E deficiency manifests in the early stage of atherosclerosis and might be the most sensitive parameter for this.

Measures that decrease prooxidant stimuli or enhancement of antioxidant defenses may reduce risk of clinical events from atherosclerosis, however interventional trials are required to support the role of vitamin E in this regard.

## REFERENCES

1. Porter M, Pearson DJ, Suarezmeder VJ, Blann AD. Plasma, platelet and erythrocyte glutathione peroxidases as risk factors in ischaemic heart disease in man. *Clin Sci* 1992; 83: 343-5.
2. Hill MF, Singhal PK. Antioxidant and oxidative stress changes during heart failure subsequent to myocardial infarction in rats. *Am J Pathol* 1996; 148: 291-300.

3. Ross R. The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature* 1993; 362: 801-9.
4. Steinbrecher UP, Zhang HF, Loughheed M. Role of oxidatively modified LDL in atherosclerosis. *Free Radic Biol Med* 1990; 9: 155-68.
5. Esterbauer H, Gebicki J, Puhl H, Jurgens G. The role of lipid peroxidation and antioxidants in oxidative modification of LDL. *Free Radic Biol Med* 1992; 13: 341-90.
6. Witztum JL. The oxidation hypothesis of atherosclerosis. *Lancet* 1994; 344: 793-5.
7. Gey KF. Ten-year retrospective on the antioxidant hypothesis of atherosclerosis: threshold plasma levels of antioxidant nutrients related to minimum cardiovascular risk. *J Nutr Biochem* 1995; 6:206-36.
8. Simon E, Paul JL, Soni T, Simon A, Moatt N. Plasma and erythrocyte vitamin E content in asymptomatic hypercholesterolemic subjects. *Clin Chem* 1997; 43: 285-9.
9. Jayakumari N, Ambika Kumari V, Balakrishnan KG, Subramaniam Iyer K. Antioxidant status in relation to free radical production during stable and unstable anginal syndromes. *Atherosclerosis* 1992; 94: 183-90.
10. Placer ZA, Cushmann LL, Johnson BC. Estimation of product of lipid peroxidation (malonyldialdehyde) in biological systems. *Ann Biochem* 1996; 16: 259-64.
11. Alanson LG, Warwick WJ. A fluorometric micro-method for serum tocopherol. *Am J Clin Path* 1996; 36: 133-8.
12. Misra HP, Fridovich I. The role of superoxide anion in the autooxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem* 1972; 247: 3170-5.
13. Beutler E, Duron O, Kelly BM. Improved method for the determination of blood glutathione. *J Lab Clin Med* 1963; 61: 882-8.
14. Keane JF, Simon DI, Freedman JE. Vitamin E and vascular homeostasis: implications for atherosclerosis. *FASEB J* 1999; 13: 965-76.
15. Keane JF, Gou Y, Cunningham D, Shwaery GT, Xin A, Vita JA. Vascular incorporation of alpha-tocopherol prevents endothelial dysfunction due to oxidized LDL by inhibiting protein kinase C stimulation. *J Clin Invest* 1996; 98: 386-94.
16. Boscoboinik D, Szenczyk A, Azzi A. Alpha-tocopherol (vitamin E) regulates vascular smooth muscle cell proliferation and protein kinase C activity. *Arch Biochem Biophys* 1991; 186: 264-9.
17. Oury TD, Day BJ, Crapo JD. Extracellular superoxide dismutase: A regulator of nitric acid bioavailability. *Lab Invest* 1996; 75: 617-36.

**Corresponding author:**

Dr. Simmi Kharb  
1447, Sector 1, Urban Estate  
Rohtak - 124001  
Haryana, India  
E-mail: [simmikh@rediffmail.com](mailto:simmikh@rediffmail.com)